

VERTICAL DISTRIBUTION OF ANTARCTIC SOIL ALGAE  
BY DIRECT OBSERVATION WITH THE CONTACT  
SLIDE METHOD (EXTENDED ABSTRACT)

Masaru AKIYAMA<sup>1</sup>, Shuji OHTANI<sup>2</sup> and Hiroshi KANDA<sup>2</sup>

<sup>1</sup>Department of Biology, Faculty of Education, Shimane University,  
Nishikawatsu, Matsue 690

<sup>2</sup>National Institute of Polar Research, 9-10, Kaga 1-chome, Itabashi-ku, Tokyo 173

The dilution method has been commonly adopted for the counting of soil microbes, and the method has also been adopted for many soil algal studies in the Antarctic (CAMERON *et al.*, 1968; CAMERON and DEVANEY, 1970; BROADY, 1979a, b). Recently, the chlorophyll content of soil has frequently been applied to estimating the algal biomass as biochemicals (HOSHIAI and MATSUDA, 1979; FRIEDMANN *et al.*, 1980; BROADY, 1986; AKIYAMA *et al.*, 1986). However, these methods are not always adequate for studying soil algal ecology in the survey of vertical micro-distribution of algae under the Antarctic soil condition *in situ*. On the other hand, the contact slide method designed by COLODNY (1930, cited from MATSUGUCHI, 1975) has been known as a convenient method for direct counting of soil microbes under the condition *in situ*, and it was already adopted for Antarctic soil bacteriological study (VISHNIAC and MAINZER, 1972). However, little is known of the study of Antarctic soil algae using the method. In an attempt to obtain the details particularly of vertical micro-distribution of algae, the contact slide method was tried in this study.

Frosted slide glasses (*ca.* 76×26mm) were set up in January 1988 at three sites along a stream in the Yukidori Valley, Langhovde, Antarctica, and they were collected in January 1989. At each site, slide glasses were buried at a few meters intervals from the stream shore along the line set at a right angle to the stream.

The occurrence of soil algae on the surface of each slide glass was recognized between the top portion of most of the slide glasses which corresponded to the soil surface and the lower portion of the slide glass which corresponded to the soil of 5cm depth. Eleven taxa of Cyanophyceae, eight taxa of Chlorophyceae and four taxa of Bacillariophyceae were recognized (Table 1). Among them, *Actinotaenium cucurbita* and *Pinnularia borealis* dominantly occurred on the surface of the slide glasses at all of the researched sites. The algal cell numbers of dominant species ranged from 0 to 300 cells/mm<sup>2</sup>. These values do not always correspond directly to the values of soil algal biomass obtained by means of dilution method, since the values obtained by the dilution method usually give a cell quantity per volume or mass of soil, whereas the values obtained by the contact slide method show merely a cell quantity per area of vertically sectioned surface of soil. However, the values obtained by the contact slide method are closely related to the values of chlorophyll content of soil which were derived mainly from epipsamic algae at the same site ( $r=0.75$ ).

Table 1. List of algae detected from the surface of the slide glasses buried in the Antarctic sandy soil.

---

CYANOPHYCEAE

- Synechococcus aeruginosus* NÄG.  
 \**Synechococcus maior* SCHR.  
*Gloeocapsa magma* (BRÉB.) KÜTZ.  
*Gloeocapsa ralfsiana* KÜTZ.  
*Gloeocapsa compacta* KÜTZ.  
 \**Lyngbya martensiana* MENEGH.  
*Lyngbya* sp. 1 (width=1 µm)  
*Lyngbya* sp. 2 (width=3 µm)  
*Nostoc commune* VAUCHER  
*Nostoc* sp.  
*Stigonema minutum* (AG.) HASS.

BACILLARIOPHYCEAE

- Hantzschia amphioxys* (EHR.) GRUN.  
*Navicula muticopsis* VAN HEURCK  
*Navicula arcuata* HEID. et KOLBE  
 \**Pinnularia borealis* EHRENB.

CHLOROPHYCEAE

- \**Actinotaenium cucurbita* (BRÉB.) TEILING.  
*Stichococcus* sp.  
 Coccoid alga no. 1 (diam.=10 µm; tetrads)  
 Coccoid alga no. 2 (diam.=10–16 µm; unicell)  
 Coccoid alga no. 3 (diam.=6–8 µm; colonies with gelatinous envelopes)  
 Coccoid alga no. 4 (irregularly shaped colonies composed of densely aggregated cells)  
 Coccoid alga no. 5 (colonies composed of 1, 2, 4, to 16 cells with thick colonial gelatinous envelopes)  
 Filamentous alga
- 

\* indicates species dominant occurrence.

Horizontal distribution of soil algae along the stream showed a steep decrease in both chlorophyll content and cell numbers, from the water boundary area towards the site away from the stream.

The structure of algal communities, particularly their vertical micro-distribution, varied with site, viz. the depth showing a maximal algal growth at each site was as follows; –0.5 cm (53 cells/mm<sup>2</sup>) at site 1, –1 cm (318 cells/mm<sup>2</sup>) at site 2, and –2 cm (555 cells/mm<sup>2</sup>) at site 3.

Patterns of vertical distribution of each algal species varied with site, but the factors which determined their micro-distribution could not be clarified.

References

- AKIYAMA, M., OHYAMA, Y. and KANDA, H. (1986): Soil nutrient condition related to the distribution of terrestrial algae near Syowa Station, Antarctica. Mem. Natl Inst. Polar Res., Spec. Issue, 44, 198–201.  
 BROADY, P. A. (1979a): The Signy Island terrestrial reference sites; IX. The ecology of the algae of Site 2, a moss carpet. Br. Antarct. Surv. Bull., 47, 13–29.

- BROADY, P. A. (1979b): Quantitative studies on the terrestrial algae of Signy Island, South Orkney Islands. *Br. Antarct. Surv. Bull.*, **47**, 31–41.
- BROADY, P. A. (1986): Ecology and taxonomy of the terrestrial algae of the Vestfold Hills. *Antarctic Oasis*, ed. by J. PICKARD. Sydney, Academic Press, 165–202.
- CAMERON, R. and DEVANEY, J. R. (1970): Antarctic soil algal crusts; Scanning electron and optical microscope study. *Trans. Am. Microsc. Soc.*, **89**, 264–273.
- CAMERON, R., KING, J. and DAVID, C. (1968): Soil microbial and ecological studies in southern Victoria Land. *Antarct. J. U.S.*, **3**, 121–123.
- FRIEDMANN, E. I., LAROCK, P. A. and BRUNSON, J. O. (1980): Adenosine triphosphate (ATP), chlorophyll, and organic nitrogen in endolithic microbial communities and adjacent soils in the dry valleys of southern Victoria Land. *Antarct. J. U.S.*, **15**, 164–166.
- HOSHIAI, T. and MATSUDA, T. (1979): Aderie penguin rookeries in the Lützow-Holm Bay area and relation of rookery to algal biomass in soil. *Mem. Natl Inst. Polar Res., Spec. Issue*, **11**, 140–152.
- MATSUGUCHI, T. (1975): *Zenkinsuu no sokutei* (Measurement of soil microbes). *Dojô Biseibutsu Jikkenho* (Manual of Soil Microbial Experiments), ed. by Jpn. Soc. Soil Microbes. Tokyo, Yokendo, 27–31.
- VISHNIAC, W. V. and MAINZER, S. E. (1972): Soil microbiology studied *in situ* in the dry valleys of Antarctica. *Antarct. J. U.S.*, **7**, 88–89.

(Received May 17, 1990; Revised manuscript received July 18, 1990)